



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ASTRONOMICAL OBSERVATIONS IN 1908.

MADE BY TORVALD KÖHL, AT ODDER, DENMARK.

VARIABLE STARS.

(The instrument used is a 3-inch Steinheil, power 42.)

S Ursæ Majoris.¹

Jan.	2:	S { $> e.$ $< d.$	Apr.	8:	id.
	4:	id.		19:	1 step $> f.$
	5:	$< d.$	May	1:	= f.
	10:	1 step $< d.$		11:	= g.
	13:	= d.		18:	{ $< g.$ $> h.$
	14:	id.		20:	id.
	18:	id.	Aug.	14:	1 step $> d.$
	20:	id.		17:	2 steps $> d.$
	21:	id.		23:	id.
	28:	id.		24:	id.
	29:	id.		27:	id.
	31:	id.		29:	1 step $> d.$
Feb.	2:	1 step $> d.$		31:	2½ steps $> d.$
	6:	2 steps $> d.$	Sept.	3:	3 steps $> d.$
	8:	id.		9:	id.
	10:	id.		16:	5 steps $> d.$
	16:	id.		19:	6 steps $< c.$
	19:	{ in the midst between d and c.		22:	5 steps $< c.$
	23:	2½ steps $> d.$		24:	id.
	29:	{ in the midst between d and e.		25:	id.
Mar.	23:	1 step $> e.$		30:	5 steps $> d.$
	24:	= e.	Oct.	3:	id.
	25:	1 step $< e.$		5:	id.
	26:	id.		10:	4 steps $> d.$
	27:	id.		18:	1 step $< d.$
	28:	2 steps $< e.$		22:	2 steps $> d.$
	29:	id.	Nov.	1:	= e.
	30:	3 steps $< e.$		6:	id.
Apr.	1:	4 steps $< e.$		8:	id.
	2:	3 steps $> f.$		16:	{ in the midst between e and f'.
	3:	id.		26:	2 steps $> f.$
	6:	id.		29:	{ $< f.$ $> g.$
	7:	id.			

¹ Vide the sketch in the *Publications A. S. P.*, No. 73, 12, 56.

*T Ursæ Majoris.*¹

Jan. 2:	T 5 steps > a.	Apr. 8:	invisible.
4:	4 steps > a.	15:	id.
5:	id.	16:	id.
10:	2 steps > a.	19:	id.
13:	1 step > a.	20:	id.
14:	id.	May 11:	id.
18:	2 steps > a.	18:	id.
20:	1½ step > a.	20:	id.
21:	2 steps > a.	Aug. 14:	{ < b.
28:	1 step < a.	17:	{ > c.
29:	id.	23:	{ 1 step > b.
31:	id.	24:	{ < a.
Feb. 2:	2 steps < a.	27:	{ 2 steps > b.
6:	{ < b.	29:	{ < a.
8:	{ > c.	31:	{ > b.
9:	1 step < b.	Sept. 3:	{ 1 step > a.
10:	id.	9:	{ 2 steps > a.
16:	2 steps > c.	16:	{ 3 steps > a.
19:	id.	19:	{ id.
19:	{ in the midst	22:	{ id.
23:	{ between c and d.	24:	{ id.
23:	1½ step > d.	25:	{ id.
29:	{ in the midst	30:	{ = a.
29:	{ between d and e.	Oct. 3:	{ id.
Mar. 18:	< e.	5:	{ = b.
23:	= f.	10:	{ id.
24:	{ < f.	18:	{ = c. (?)
25:	{ > g.	22:	{ < c.
26:	id.	Nov. 1:	{ > d.
27:	1 step > g.	4:	{ 1 step < d.
28:	id.	6:	{ 3 steps > e.
29:	1 step < g.	8:	{ id.
30:	id.	16:	{ id.
Apr. 1:	very faint.	26:	{ = g.
2:	id.	29:	{ = f = g.
3:	id.		
6:	id.		
7:	id.		

The comparison stars f and g are found to be a little variable. August 24th, I have noted: g 1 step > f. On November 29th: f = g. Usually I note: f > g, the difference being only 1 step. B. D. has f = g = 9^m.5; Harvard has f = 10^m.75, g = 10^m.40.

¹ Vide the sketch in the *Publications A. S. P.*, No. 22, 4, 63.

*W Pegasi.*¹

Jan.	1:	W = b.	Aug.	24:	id.
	2:	{ in the midst		27:	{ < g.
		{ between b and c.			{ > h.
	4:	{ < b.		29:	id.
		{ > c.		31:	id.
	5:	id.	Sept.	3:	id.
	14:	1 step > c.		4:	id.
	18:	= c.		12:	id.
	20:	id.		16:	a little > h.
	21:	1 step < c.		20:	{ < g.
	29:	3 steps < c.			{ > h.
	31:	1 step > d.		24:	= g.
Feb.	2:	1½ step > d.		25:	= f.
	9:	1 step > d.		30:	id.
	19:	= e.	Oct.	3:	1 step < f.
	29:	{ in the midst		5:	id.
		{ between e and f.		18:	1 step < e.
Aug.	14:	{ < g.	Nov.	8:	2 steps < c.
		{ > h.		16:	1 step < b.
	18:	= h.		26:	= b.
	23:	id.		29:	id.

*SS Cygni.*²

Jan.	1,	6 ^h :	SS = g.	May	11, 11 ^h :	{ 3 steps > c.
	2,	9 ^h :	< g.			{ 1 step < b.
	4,	6 ^h :	= h.		15, 10 ^h :	1 step > c.
	5,	7 ^h :	very faint.		18, 11 ^h :	1 step < c.
	14,	6 ^h :	invisible.		20, 12 ^h :	= d.
	18,	6 ^h :	id.	Aug.	14, 10 ^h :	{ > e.
	20,	6 ^h :	id.			{ < d.
	21,	7 ^h :	id.		15, 10 ^h :	{ in the midst
	29,	7 ^h :	1 step < b.			{ bet. d and e.
	31,	7 ^h :	= b.		17, 11 ^h :	< e.
Feb.	2,	7 ^h :	{ < b.		18, 10 ^h :	4 steps < e.
			{ 2 steps > c.		23, 10 ^h :	= f.
	6,	6 ^h :	2 steps > d.		24, 10 ^h :	{ > f.
	9,	6 ^h :	= e.			{ < e.
	16,	7 ^h :	invisible.		27, 10 ^h :	1 step > f.
	19,	7 ^h :	very faint.		29, 10 ^h :	{ > f.
Apr.	20, 12 ^h :	invisible.				{ < e.
	21, 10 ^h :	id.			31, 10 ^h :	= e.
	26, 11 ^h :	id.		Sept.	3, 10 ^h :	= e.
May	1, 11 ^h :	1 step < e.			4, 9 ^h :	{ > f.
	9, 10 ^h :	1 step < c.				{ < e.

¹ Vide the sketch in the *Publications A. S. P.*, No. 60, 10, 23.

² Vide the sketch in the *Publications A. S. P.*, No. 100, 17, 18.

Sept. 9, 9 ^h : = d.	Oct. 5, 8 ^h : { < d'. > e.
11, 8 ^h : 2 steps < c.	18, 9 ^h : d'(3)SS(2)e
12, 8 ^h : 1½ step < c.	Nov. 1, 8 ^h : { < d'. > e.
16, 9 ^h : 2 steps > c.	4, 8 ^h : = e.
19, 9 ^h : { in the midst	6, 9 ^h : = e(?).
20, 8 ^h : b(3)SS(2)c	8, 7 ^h : d'(3)SS(2)e
23, 8 ^h : { < c. > d.	14, 7 ^h : 2 steps > c.
24, 10 ^h : = d.	16, 8 ^h : = d.
25, 9 ^h : 2 steps < d.	26, 6 ^h : = e.
30, 12 ^h : 4 steps < d.	29, 6 ^h : 1 step > f.
Oct. 1, 10 ^h : id.	
3, 9 ^h : = e.	

Z Cygni.¹

Jan. 1: Z 2 steps > a.	Aug. 14: id.
2: 1 step > a.	18: id.
4: < a.	23: id.
5: { < a. > b.	Sept. 3: = e.
14: 1 step > a.	9: { > e. < d.
18: 2 steps > a.	11: { > d. < c.
20: 1 step > a.	16: id.
21: id.	19: = d.
28: = a.	22: { > c. < b.
29: 2 steps > a.	24: id.
31: 1 step > a.	30: = b.
Feb. 2: = a.	Oct. 3: = b'.
6: = b.	5: id.
16: 2 steps > b(?).	18: { < a. > b'.
19: = b.	Nov. 1: = b'.
Mar. 25: very faint.	4: = b.
Apr. 1: id.	6: 1 step > b.
20: invisible.	8: 1½ step < b.
21: id.	14: 1 step < b.
26: very faint.	16: { < b. > c.
May 1: faint, < e.	29: = d.
9: invisible.	
15: id.	
18: id.	
20: id.	

Y Tauri (B. D. + 20° 1083).

A number of forty comparisons have been made upon this irregular variable star, which during the whole year was either

¹ Vide the sketch in the *Publications A. S. P.*, No. 100, 17, 16.

equal to or some steps brighter than the star $A = B. D. + 20^{\circ} 1095$ ($7^m.4$). In February, March, and April the star Y had reached its maximum ($7^m.1$), but in January and November it had decreased a few steps.

U Herculis.

Apr. 20: U invisible(?).	Sept. 11: $\left\{ \begin{array}{l} < c. \\ > d. \end{array} \right.$
26: 2 steps < h.	12: = c.
May 1: id.	16: $\left\{ \begin{array}{l} < c. \\ > d. \end{array} \right.$
18: 1 step > h.	19: C(2)U(3)d.
20: id.	22: id.
Aug. 15: = a.	23: $\left\{ \begin{array}{l} \text{in the midst} \\ \text{between c and d.} \end{array} \right.$
17: $\left\{ \begin{array}{l} > c \\ < a \end{array} \right\} b > a.$	24: id.
18: 2 steps < a.	25: C(3)U(2)d.
23: id.	30: id.
24: id.	Oct. 3: = d.
27: id.	5: 2 steps > d.
29: id.	10: 1 step > d.
31: = c.	18: = e.
Sept. 3: id.	Nov. 29: 2 steps < f.
4: 1 step > c.	
9: $\frac{1}{2}$ step > c.	

I have used the sketch in the *Publications A. S. P.*, No. 106, 18, 52, drawn by Miss ROSE O'HALLORAN, but have added the two small neighboring stars g at a and h at f, both northward.

A SUSPECTED VARIABLE STAR.

138.1908 *Herculis*.

During my observations upon *U Herculis* my attention was directed to the two comparison-stars,

$$\begin{aligned} a &= B. D. + 19^{\circ} 3096 & 7^m.0. \\ b &= B. D. + 19^{\circ} 3089 & 7^m.8. \end{aligned}$$

The star b is here eight steps fainter than a. On August 5, 1907, I found $b > a$, and so I have seen it until October, 1908. On October 5th, 10th, and 18th, I have noted: b 2-3 steps < a. But when the ocular was screwed out the reddish star a was "diminished," and then seemed to be equal to or even dimmer than b. In A. G. Berlin A 5856 is

$$\begin{aligned} a &= 7^m.2 \\ b &= 7^m.7, \end{aligned}$$

and in *Harvard Annals*, 37, 170 and 183, we find

$$a = 7^m.36$$

$$b = 7^m.39.$$

Whether the variable is a or b is still difficult to decide. According to its red color, it might perhaps be the star a, though a look at the differences might point out b as the variable one. It will be necessary to compare b with a and c. (Vide *Astronomische Nachrichten*, No. 4274, 179, 29.)

METEORS.

Fireballs have been observed from stations in Denmark at the following dates: January 3d, 21st; March 2d, 16th, 25th; April 8th, 15th; May 1st, 11th, 29th; June 25th, 26th; July 2d, 20th, 21st; August 6th, 20th; October 7th, 24th; November 8th, 21st; December 3d, 7th.

SHOOTING-STARS.

A little swarm of shooting-stars, during five minutes more than thirty meteors, was observed on January 2d, 8^h 10^m–8^h 20^m, at Hjörning, Denmark, radiant = 300° + 61°; and on January 3d, 11^h 23^m–12^h 42^m, seventeen large shooting-stars were mapped at Paderborn, Germany. (Vide *Astronomische Rundschau*, No. 98, and *Astronomische Nachrichten*, No. 4263, 178, 255.) On June 25th, 10^h 30^m, an observer at Odense, Denmark, mapped a most interesting twin-meteor, two shooting-stars with parallel paths, with a distance of ½°, one on each side of the pretty star *Arcturus*.

The weather was, in the year 1908, quite unfavorable for the planned observations on shooting-stars in August, and after November 29th the sky was overcast with clouds every night here at Odder.

The above-mentioned estimations of variable stars have often been controlled by my young assistant, JÖRGEN FOG.